CS457 - Absolute Discount Smoothing Handout

David Kauchak - Fall 2011

Given the following corpus (where we only have one letter words):

aaab

a b b a

caaa

We would like to calculate an absolute discounted model with D = 0.5. We'll ignore the begin and end sentence tokens as well and assume that our vocabulary is all three "words".

We first calculate the unigram MLE probabilities as:

	MLE prob
p(a)	8/12
p(b)	3/12
p(c)	1/12

and the bigram MLE probabilities as:

	MLE prob	
p(a a)	4/6	
p(b a)	2/6	
p(c a)	0	
p(a b)	1/2	
p(b b)	1/2	
p(c b)	0	
p(a c)	1	
p(b c)	0	
p(c c)	0	

Using this information, we can now calculate the reserved mass and the α s for each of the words in our vocabulary (i.e. things that we're conditioning on):

• a

The numerator for our α is the reserved mass:

$$reserved_mass(a) = (2 * 0.5)/6 = 1/6$$

and the denominator is:

$$1 - \sum_{x:count(ax)>0} p(x) = 1 - (p(a) + p(b)) = 1 - (8/12 + 3/12) = 1/12$$

giving us an α of:

$$\alpha(a) = \frac{1/6}{1/12} = 2$$

• *b*

$$reserved_mass(b) = (2*0.5)/2 = 1/2$$

$$1 - \sum_{x:count(bx)>0} p(x) = 1 - (p(a) + p(b)) = 1 - (8/12 + 3/12) = 1/12$$

$$\alpha(b) = \frac{1/2}{1/12} = 6$$

• (

$$reserved_mass(c) = (1*0.5)/1 = 1/2$$

$$1 - \sum_{x:count(bx)>0} p(x) = 1 - p(a) = 1 - (8/12) = 4/12 = 1/3$$

$$\alpha(c) = \frac{1/2}{1/3} = 3/2$$

Finally, now that we have the α s, we can calculate the smoothed bigram probabilities. For those that occurred, we simply discount the count. For those that did not occur, we calculate the probability as alpha times the unigram probability of the word.

	eqn	prob
p(a a)	(4 - 0.5)/6	3.5/6
p(b a)	(2 - 0.5)/6	1.5/6
p(c a)	2 * 1/12	1/6
p(a b)	(105)/2	1/4
p(b b)	(1-0.5)/2	1/4
p(c b)	6 * 1/12	1/2
p(a c)	(1-0.5)/1	1/2
p(b c)	3/2 * 3/12	3/8
p(c c)	3/2 * 1/12	1/8

Notice that after the smoothing, the three distributions all still sum to 1. In this case discounting by 0.5 may be a bit aggressive.