
csci54 – discrete math & functional programming
proofs on sets, functions

-
- ▶ this week:
 - ▶ week09-group
 - ▶ week09-ps
 - ▶ next week:
 - ▶ Tuesday review session (based on week09-group)
 - ▶ Thursday checkpoint
 - ▶ open one 8.5"x11" double-sided page of notes
 - ▶ closed everything else
 - ▶ Focused on things covered after the first checkpoint
 - ▶ no group meeting (or optional, reach out to your TA), no problem set
 - ▶ week after that:
 - ▶ Number theory, final bits of 54
-



Recall: sets

- ▶ A set is an unordered collection of objects
- ▶ Given a set S and an object o , either $o \in S$ or $o \notin S$
- ▶ The cardinality of a set is written $|S|$ and is the number of elements in the set
- ▶ Special sets:
 - ▶ the empty set, which contains no elements: $\{\}$ or
 - ▶ the universal set, U
- ▶ Set operations: complement, union, intersection, difference, Cartesian product
- ▶ Comparing/relating sets: equality, subset, proper subset, superset, proper superset, disjoint



Set operations

- ▶ S^c : set complement
 - ▶ $S^c = \{ x \in U : x \notin S \}$
- ▶ $S \cup T$: set union
 - ▶ $S \cup T = \{ x : x \in S \text{ or } x \in T \}$
- ▶ $S \cap T$: set intersection
 - ▶ $S \cap T = \{ x : x \in S \text{ and } x \in T \}$
- ▶ $S - T$: set difference
 - ▶ $S - T = \{ x : x \in S \text{ and } x \notin T \}$
- ▶ $S \times T$: Cartesian product
 - ▶ $A \times B = \{ (x, y) : x \in A \text{ and } y \in B \}$

Comparing/relating sets

- ▶ $=$: set equality
- ▶ S and T contain the same elements
- ▶ \subseteq : subset
- ▶ S contains T
- ▶ \subset : proper subset
- ▶ S contains T and S does not equal T
- ▶ \supseteq : superset
- ▶ T contains S
- ▶ \supset : proper superset
- ▶ T contains S and T does not equal S
- ▶ “ T and S are disjoint”
- ▶ T and S share no elements



proofs on sets

- ▶ Element-wise:

- 📖 Show that no matter which elements of the sets are picked, membership/non-membership is provable

- ▶ Algebraic:

- 📖 Use properties of the operations to show relations between sets



proofs on sets

- ▶ Claim $\{x \in \mathbb{Z} : 18|x\} \subseteq \{x \in \mathbb{Z} : 6|x\}$

let y be an arbitrary element of the set $\{x \in \mathbb{Z} : 18|x\}$
then direct proof using definition of subset

- ▶ Proof:
 - ▶ Let y be an arbitrary element of the set $\{x \in \mathbb{Z} : 18|x\}$
 - ▶ This means there exists an integer k such that $y = 18k$
 - ▶ Furthermore, since $y = 18k = 6(3k)$, we know that $6|y$ and $y \in \{x \in \mathbb{Z} : 6|x\}$
 - ▶ Since this is true for any element $y \in \{x \in \mathbb{Z} : 18|x\}$, it is true for every element.
 - ▶ Therefore $\{x \in \mathbb{Z} : 18|x\} \subseteq \{x \in \mathbb{Z} : 6|x\}$



proofs on sets

- ▶ Claim $\{x \in \mathbb{Z} : 18|x\} \subseteq \{x \in \mathbb{Z} : 6|x\}$
 - ▶ Is the claim still true if we replace superset with strict superset? Prove your answer correct
 - ▶ Is the claim still true if we replace superset with equals? Prove your answer correct
 - ▶ More generally:
 - ▶ how do you prove set equality?
 - ▶ prove subset in both directions
 - ▶ how do you prove strict subset?
 - ▶ prove subset and not equal
-






Recall: functions

Definition 2.46: Function.


Let A and B be sets. A *function* f from A to B , written $f : A \rightarrow B$, assigns to each input value $a \in A$ a unique output value $b \in B$; the unique value b assigned to a is denoted by $f(a)$. We sometimes say that f *maps* a to $f(a)$.

- ▶ Given a function
 - ▶ the domain is the set A
 - ▶ the co-domain is the set B
 - ▶ the range (or the image) is the subset of B that are actually mapped to by an element in A .




classifying functions - definitions

- ▶ one-to-one: a function is one-to-one if, for every element of the co-domain, at most one element of the domain maps to it.
- ▶ onto: a function is onto if, for every element of the co-domain, there is an element of the domain that maps to it.
 - ▶ alternatively, a function is onto if the co-domain equals the range
- ▶ bijection: a function is a bijection if it is both one-to-one and onto



classifying functions

- ▶ one-to-one: a function is one-to-one if, for every element of the co-domain, at most one element of the domain maps to it.
 - ▶ in other words: if $f(x) = f(y)$, then $x=y$.
- ▶ onto: a function is onto if, for every element of the co-domain, there is an element of the domain that maps to it.
 - ▶ in other words, $f(y) = x$



example

- ▶ Claim: the function $g(x) = x-1$ is a bijection

a function is a bijection if it is both one-to-one and onto

- ▶ Proof:

- ▶ g is one-to-one:

one-to-one: if $f(x) = f(y)$, then $x=y$

- ▶ assume there are two elements x, y in Z such that $g(x)=g(y)$. Then $x-1 = y-1$, so $x=y$

- ▶ therefore g is one-to-one

onto: $f(y) = x$

- ▶ g is onto:

- ▶ let x be any element of Z . Then $x+1$ is an element that maps to x .
 - ▶ since x is any element of Z , every element of Z has an element that maps to it,
 - ▶ therefore g is onto

-
- ▶ ▶ since g is one-to-one and onto, g is a bijection

a little more on bijections

- ▶ If a function is a bijection, then it is also invertible. In other words, if f is a bijection, then there is a function f^{-1} such that $f(x) = y$ iff $f^{-1}(y) = x$
- ▶ The identity function $f(x)=x$ is a bijection
 - ▶ the identity function is the function that maps every element to itself

